







- [2] Cullen J, in Davis PH. 1965. Flora of Turkey and the East Aegean Islands, Edinburgh Univ. Press, Edinburgh, volume: 1: 452-460.
- [3] Davis PH, Mill RR, Tan K. (eds), 1988. Flora of Turkey and the East Aegean Islands (supplement), Edinburgh, volume: 10: 50-54.
- [4] Güner A, Özhatay N, Ekim T, Başer KHC. (eds), 2000, Flora of Turkey and the East Aegean Islands (supplement), Edinburgh, volume: 11.
- [5] Seçmen Ö, Gemici Y, Görk G, Bekat L, Leblebici E. 1998. Tohumlu Bitkiler Sistematigi, Ege Üniv. Basımevi, Bornova-İzmir, 236-240.
- [6] Şimşek I, Aytekin F, Yeşilada E, Yıldırım Ş. 2002. Anadolu'da Halk Arasında Bitkilerin Kullanılış Amaçları Üzerinde Etnobotanik Bir Çalışma, 14. Bitkisel İlaç Hammaddeleri Toplantısı, Eskişehir.
- [7] Blackwell WH. 1991. Poisonous and Medicinal Plants, Systematic Botany, 16(1): 206-207.
- [8] Martin E, Duran A, Ocak A, Dinç M, Öztürk M. 2008. A Cytotaxonomical Study on Endemic *Hesperis turkmenaghensis* A.Duran & A.Ocak (Sect. *Hesperis*) (Cruciferae), IVth Balkan Botanical Congress, Sofia, Phytologia Balcanica, Bulgaria.
- [9] Ünal F, Duran A, Martin E. 2008. Karyotype on the genus *Hesperis* L. (Cruciferae) of the *Diaplectos* (Dvořák) Dvořák section in Turkey, Hereditas, 145: 64-68.
- [10] Kochjarová J, Valachovi M, Bure P, Mráz P. 2006. The genus *Cochlearia* L. (Brassicaceae) in the Eastern Carpathians and adjacent area, Botanical Journal of the Linnean Society, 151: 355-364.
- [11] Kolník M, Marhold K. 2006. Distribution, chromosome numbers and nomenclature conspect of *Arabidopsis halleri* (Brassicaceae) in the Carpathians, Biologia, 61: 141-50
- [12] Olin-Fatih M. 1996. The morphology, cytology, and C-banded karyotypes of *Brassica campestris*, *B. oleracea*, and *B. napus* plants regenerated from protoplasts. TAG Theoretical and Applied Genetics, 93(3): 414-420.
- [13] Grundt H, Obermayer R, Borgen L. 2005. Ploidal levels in the arctic-alpine polyploid *Draba lactea* (Brassicaceae) and its low-ploid relatives, Botanical Journal of the Linnean Society 147(3): 333-347.
- [14] Mukherjee P. 1975. Karyomorphological studies of ten strains of Indian mustard (*Brassica juncea* Coss.), Euphytica, 24(2): 483-486.
- [15] Marhold K, Ancëv EM, Tan K. 1996. A new subspecies of *Cardamine amara* (Brassicaceae) from Bulgaria and Greece. Ann. Bot. Fennici, 33: 199-204.
- [16] Tsukaya H, Yokoyama J, Ikeda H, Kuroiwa H, Kuroiwa T, Iwatsuki K. 1997. Morphological, Physiological and Molecular Genetic Characterization of *Arabidopsis himalaica*, J. Plant Res., 110: 15-23.